

CHAMBERS, S. 2009.

Birds of New Zealand - Locality Guide. 3rd edn.

Arun Books, Orewa, New Zealand.

Antarctic fulmar: p 50.



### ANTARCTIC FULMAR

**Family** Procellariidae

**Species** *Fulmarus glacialoides*

**Common names** Antarctic Fulmar, Silver-grey Fulmar

**Status** Native

**Abundance** Uncommon.

**Size** 500 mm (House Sparrow 145 mm, Flesh-footed Shearwater 450 mm)

**Breeding islands close to New Zealand** On islands at the fringes of the Antarctic.

**Breeding months** December onwards in colonies. One white egg is laid in a shallow scrape.

**Range worldwide** Around southern oceans.

**New Zealand range** Widely spread north to the Bay of Plenty on the east coast and to the Kaipara Harbour on the west. At Kaikoura can be seen feeding close to land especially when the weather is rough.

## **Description**

**Head, neck and underparts** White. **Underwing** White with black trim to trailing edges. **Upperwings and mantle** Soft grey, with dark edges to primary feathers.

**Tail** White. **Bill** Pinkish with dark tip. **Legs and feet** Pinkish blue.

## **Conspicuous features**

- Larger than Flesh-footed Shearwater.
- White head and tail.

## **Conspicuous characteristics**

- Usually seen alone.
- Will join in seabird feeding frenzies.

## **Where to find – South Island**

**Kaikoura** – From pelagic excursions. Also seen from Stewart Island.