

A large diverse group of birds of estuaries, coasts, riverbeds and farmland. Most are long-legged and feed in or near shallow water. Bill shape is varied; short and stubby in those (e.g. dotterels) that peck from the surface, but longer in those that feed in shallow water (e.g. stilts), or probe deeply (e.g. godwits). Flight strong and direct. Often form flocks while roosting or flying, but disperse to feed. Many species seen in NZ breed in the Arctic and arrive in September, with remnants of breeding plumage, and depart in March, often in breeding plumage. Most subadults and a few adults spend the southern winter here.

COMMON SANDPIPER *Tringa hypoleucos*

Rare Arctic migrant

20 cm, 50 g. Small dark solitary sandpiper with *white tab in front of folded wing*. Often perches on rocks or logs and bobs head and tips tail. Jerky, flickering flight interspersed with glides on stiffly held wings. Head and upperparts olive brown except white eye-ring and eyebrow, boldest in front of eye; throat and sides of neck washed brown; underparts white. Tail long, extending well beyond folded wings. Short straight dark brown bill; short greenish-grey legs. In flight, white wingbar and white sides to brown rump and tail. Flight call a distinctive clear 'hee-dee-dee'. **Habitat:** Breeds Eurasia. A few reach NZ, usually in small muddy estuaries. [Sp 220]



SNIPE, SANDPIPERS, GODWITS and CURLEWS

Scolopacidae

About 79 species, of which 2 breed in New Zealand (non-migratory snipe) and 32 reach New Zealand as migrants for the northern winter.

Apart from stints (known as peeps in North America), these waders have a slender bill that is as long as or longer than the head. In curlews and godwits, the bill is sensitive and flexible at the tip, and the mandibles open during probing in soft mud or shallow water. All have long, pointed wings, rapid flight, a long neck, long legs and, for birds, a short tail. Their eyes are smaller than those of plovers and dotterels, as suits their more tactile, less visual, feeding. Gregarious when not breeding.

Their food has not been studied in detail in New Zealand. They take a variety of mudflat-burrowing crabs, small amphipod and ostracod crustaceans, polychaete worms and small gastropod and bivalve molluscs. Their diet includes insect larvae and pupae of craneflies (Tipulidae), midges (Chironomidae), beetles and flies; they may also take spiders and earthworms. They swallow grass, sedge and rush seeds, but whether by accident or design is not known for certain. The various species probe to different depths according to the length of their bill. Most of the rapid, vigorous, sewing-machine probing of medium and smaller waders is exploratory, as often is the slower, more careful probing of godwits, curlews or whimbrels.

The waders that migrate to New Zealand mostly breed in the arctic or subarctic tundra of the Northern Hemisphere and are strongly migratory. Those that breed furthest north tend to migrate furthest into the Southern Hemisphere, from the largest curlews to the smallest stints. The routes taken by the various species of wader are being elucidated by an extensive co-operative programme of banding and leg-flagging throughout the East Asian flyway. The New Zealand Wader Study

Group have caught over 4000 Arctic migrants in the Auckland area, and some of these have been marked, to enable resightings, with white plastic leg flags.

The first Arctic migrants arrive in September-October, but others trickle into the country through November. Most leave in March-April, but a variable proportion of the summer population stays behind to spend the southern winter in New Zealand; most are probably yearlings, as few adopt breeding plumage, and so the number left behind provides an indication of the success of the previous northern breeding season.

New Zealand is at the southern limit of many species, and some of the distances travelled are huge; it is possible that some of the migrants fly between New Zealand and China, Japan or the Aleutian Chain in a single flight, although most stop at least once to refuel. In order to undertake such a long journey, waders feed voraciously in the weeks before departure and often arrive late to their roosts. They lay down extensive fat deposits, their weight can increase to 50-75% above their normal non-breeding weight. On arrival, they are often exhausted and quite approachable, but quickly regain their condition. The adult Arctic migrants moult all their flight feathers during the southern summer, and so can be distinguished from juveniles, which do not moult or lose only a few feathers until the southern autumn.

The sexes are alike, but females are often noticeably larger than males. The non-breeding plumage, as is mostly seen in New Zealand, is dull, the upperparts mottled or a uniform brown and grey, the underparts paler, sometimes with streaks and spots. Before leaving and while breeding, they are brighter, many species becoming much more rufous above and rufous or black below. The age of first breeding is 1-2-3 years, and many birds live to at least 15 years old.

220. COMMON SANDPIPER *Tringa hypoleucos*

Plate 45

Size: 20 cm, 50 g
Distribution: Breed across Europe and Asia from Spain and Britain to the Kamchatka Peninsula and Japan. They winter on the coasts of Africa, Madagascar, southern and southeastern Asia, the Philippines, and Australasia, especially in New Guinea. They are an uncommon visitor to New Zealand, with less than 20 sight records since they were first recorded in 1964. Only four sightings have been in the South Island: Lake Wainono

(1980), Whanganui Inlet (1981-83), Waipara Estuary (1992) and Farewell Spit (1992).
Behaviour: Usually solitary and often in places not frequented by other waders; the bird in Whanganui Inlet was by itself for three years, feeding on the same stretch of muddy tidal creek apart from other species. Common Sandpipers bob their heads, teeter up and down with their tail and fly on flickering downcurved wings.