CHAMBERS, S. 2009. Birds of New Zealand - Locality Guide. 3rd edn. Arun Books, Orewa, New Zealand. Large sand dotterel: pp 156-157.



LARGE SAND DOTTEREL

Family Charadriidae
Species Charadrius leschenaulti
Common names Large Sand Dotterel, Greater Sand Plover
Status Vagrant
Abundance Uncommon
Size 240 mm (cf sparrow 145 mm, Banded Dotterel 200mm)
Habitat A bird of estuaries and harbours.
Worldwide range Breeds in highland areas in a belt from the Caspian Sea through to Mongolia. It winters around the coasts of East Africa, southern Asia, through Malaysia and Indonesia to Australia. A few birds come to New Zealand.
New Zealand range Likely to be found in the traditional wader haunts of the northern harbours, but can be expected from any estuary.

Description – non breeding plumage

Forehead White extending to above eye. **Upperparts including shoulder tabs** Grey-brown. **Underparts** White. **Bill and legs** Black.

Description – breeding plumage

Crown nape and breast Reddish-orange. **Forehead** White, edged with black. **Eye-stripe** A black marking extends from bill through eye to back of head. **Upperparts** Brownish. **Chin** White. **Underparts** White. **Eye** Black.

Conspicuous feature

• Look for the dark eye.

Conspicuous characteristics

- Has a tendency to stand motionless.
- Runs and pauses like all plovers when feeding.

Differences between Large Sand Dotterels and Mongolian Dotterels

- Large Sand Dotterels appear leggy and stand taller than Mongolian.
- Large Sand Dotterels have a more horizontal posture.
- Large Sand Dotterels have white on chin and throat lacks black edge.
- Breast-band in breeding plumage has more orange than the Mongolian.

Where to find – North Island

North Auckland – Kaipara Harbour at Manukapua Island. P. 287.

South Auckland – Miranda. P. 302.

Where to find – South Island

Canterbury – Lake Ellesmere. P. 351.

Southland – Waituna Lagoon. P. 364.